

STUDENT'S

Pocket Prescriber

BOOKSELLER,
26, WARDOUR STREET, .
OXFORD STREET, W.

LIVINGSTONES' MEDICAL HAND-BOOKS.

EDINBURGH MEDICAL SCHOOL CALENDAR	1/3
GREVILLE'S CHEMISTRY, WITH TABLES .	9/0
MILNE'S MIDWIFERY	12/6
HUSBAND'S (AUBREY) FORENSIC MEDICINE } AND PUBLIC HEALTH	10/6
HUSBAND'S (A.) PRACTICE OF MEDICINE	7/6
HUSBAND'S (A.) POCKET PRESCRIBER .	1/0
DITTO, IN TUCK, INTERLEAVED	3/6
HUSBAND'S (A.) SANITARY LAW - .	3/6
" EXAMINATION OF THE URINE	1/0
HUGHES' MOTOR NERVES	1/0
CROOM'S MINOR GYNECOLOGY	6/0
CRAIG'S MATERIA MEDICA	9/0
CRAIG'S POSOLOGICAL TABLES . . .	1/0
CRAIG'S (BRUNN) INTERSTITIAL NEPHRITIS	4/6



22102272107

Med

ster,

K14093

THE STUDENT'S Pocket Prescriber.

BY

H. AUBREY HUSBAND, M.B.,
F.R.C.S.E.,

One of the Medical Officers to the Royal Dispensary,
Edinburgh;

Lecturer on Medical Jurisprudence and Public
Health in the Extra-Mural School.

AUTHOR OF THE "STUDENT'S HAND-BOOK OF THE
PRACTICE OF MEDICINE,"

"STUDENT'S HAND-BOOK OF FORENSIC MEDICINE
AND MEDICAL POLICE," ETC., ETC.

Edinburgh :

E. & S. LIVINGSTONE, 15 TEVIOT PLACE.

LONDON { BAILLIERE, TYNDALL, & COX ;
 { SIMPKIN, MARSHALL, & CO.

DUBLIN—FANNIN & CO. ; GILL & SON.

GLASGOW—STENHOUSE ; HADDEN ; SIME.

ABERDEEN—J. G. BISSET.

MANCHESTER—J. E. CORNISH.

1882.



309422/27569

EDINBURGH:

PRINTED FOR E. AND S. LIVINGSTONE,
BY J. T. PEDDIE AND CO.,

WELLCOME INSTITUTE LIBRARY	
Coll.	WELCOMec
Call	
No.	QV

3126-22

P R E F A C E.

THE difficulty experienced by my Pupils at the Royal Dispensary in prescribing for their out-door Patients, has induced me to prepare the following collection of Prescriptions. In so doing, I have tried to make it as comprehensive as possible, and have introduced a large number of Prescriptions for Children, in whose cases most young Prescribers find considerable difficulty.

27 CHALMERS STREET,

May 1832.

INTRODUCTION.

IN prescribing medicines for the sick, several important points have to be considered. The age, sex, temperaments, and idiosyncrasies of our patients, each and all demand our attention. Besides, we must remember that climate and disease are not without their influence, and that even the mode and time of administration are not unimportant matters to be remembered. Medicines should be made as pleasant as possible, the neglect of this simple rule often creating such a feeling of disgust on the part of the sufferer as to entirely destroy the beneficial object sought. This is especially the case with children. Let us briefly take in detail some of the points just mentioned. First, as to age. This naturally influences the dose, and for our guidance several rules have been laid down. For instance, it has been proposed to add twelve to the age, and then to divide the age by the

result. A child one year old would thus require a dose the thirteenth of that required by an adult; a child three years old, three-fifteenths, or one-fifth. Another rule equally useful is to take the age at the next birthday and divide it by twenty-four; thus the dose for one year is found by taking the age of the child at two years and dividing it by twenty-four; for example, one year $\frac{2}{24} = \frac{1}{12}$, two years $\frac{3}{24} = \frac{1}{8}$, three years $\frac{4}{24} = \frac{1}{6}$, &c. We must also not forget that children do not bear narcotics well, and that, as a rule, opium should be avoided in very young children, although they on the other hand tolerate belladonna and hyoscyamus very well. The rule for the administration of opium is one drop of the tincture for each year of age; but safe prescriptions will be given under the heading Narcotics. With regard to sex, women, as a rule, require smaller doses than men, and this is especially the case with regard to purgatives; powerful cathartics, from the straining they cause, may give rise to displacement of the pelvic organs, and, during pregnancy, to abortion. The temperaments and idiosyncrasies of individuals frequently seem to be outside the reign of all law,

and can only be known by experiment, but the effects of habit are better understood. Indulgence in alcohol, opium, and arsenic eating, renders the system so tolerant of their action, that doses poisonous when administered to others have little or no effect. The effects of climate are seen in the large doses of quiniæ that may be taken in marshy districts, and of disease in the profuse salivation produced by even small doses of mercury, in Bright's disease of the kidney, the diminished influence of narcotics in *delirium tremens* and severe pain from cancer, the passage of gall stones, &c. The risk incurred by the administration of opium in erysipelas of the head, of congestion of the brain, or of purgatives in intersusception of the bowels, are all cases in point. The question of the mode and time of administration of medicines demands more attention than is generally paid to it. The action of medicines is most rapidly produced when the drugs are injected under the skin, as they thus quickly enter the circulation, and next when they are taken upon an empty stomach. The rapid action of a purgative taken before breakfast, after the night's fast, is well

known. On the other hand, medicines which have an irritating effect on the stomach, as arsenic or iodine for instance, should be given soon after a meal. Iron appears to act best when dissolved by the acid of the gastric juice, and is, therefore, best given during active digestion. Cod liver oil also agrees best if taken with, or soon after, a meal. The action of acids in checking the secretion of the gastric juice is taken advantage of by administering them a short time before the meals in those forms of dyspepsia accompanied with excessive secretion of acid gastric juice; whereas, if the contrary effects are desired, alkalies which excite the secretion of the gastric juice should be given. The effects of valuable medicines are thus clearly influenced by their time of administration. In order that remedies may have the desired action, it is necessary that they be in a state of solution, or at least in a condition to render them more readily absorbed. Medicines, therefore, which are insoluble in water, may be readily so in the gastric juice; others again in the alkaline secretions of the pancreas and intestinal fluids. Minute division or the presence of fatty bodies

also assists the rapid absorption and action of drugs. Care must, therefore, be taken in using solutions of corrosive sublimate to the head or other parts of the body, as it has been found that poisonous effects have been rapidly produced when a solution has been allowed to spontaneously evaporate on a part. From the brief remarks just made, it will be seen that to prescribe medicines properly requires care and a due consideration of the action of each drug, and the object to be attained by its use.

In writing a prescription there are four simple rules which should be carried in the memory. In combining medicines, we seek—

1. *To augment, correct, or modify the action of a medicine.*—Thus a combination of purgatives is advisable, especially if they have each an action on different parts of the alimentary canal. Ipecacuanha increases the purgative action of jalap, and the emetic action of tartar emetic. The griping tendency of some drugs is modified by aromatics, and mercury prevented from running off by the bowels by combining it with opium.

2. *To obtain the joint operation of two or more medicines.*—The combination of purgatives just instanced will serve as an example of this rule.
3. *To obtain a new remedy.*—The union of the tincture of the perchloride of iron and the solution of the acetate of ammonia, of corrosive sublimate and the iodide of potassium, and of sulphate of iron and carbonate of potash, are all well known and useful mixtures.
4. *To afford a suitable form.*—Bread crumb and extracts for pills, emulsion of acacia to suspend heavy powders, bismuth, and the like. Glycerine disguises the taste of the tincture of iron; the essential oil of lemon that of the sulphate of magnesia.

THE STUDENT'S POCKET PRESCRIBER.

ALTERATIVES.

- ℞ Hydrargyri cum Creta, gr. iii; Pulveris Opii, gr. $\frac{1}{4}$. M. Sumat unam nocte maneque.—*In syphilis.*
- ℞ Hydrargyri Perchloridi, gr. i; Ammoniae Hydrochloratis, gr. v; Extracti Sarsæ Liquidī, ℥iiss; Decoctum Sarsæ Compositum ad ℥xii. M. Sumat ℥j ter in die.—*Constitutional syphilis, &c.*
- ℞ Hydrargyri cum Cretâ, gr. i; Pulveris Rhei, gr. i; Sodæ Bicarbonatis, gr. ii. M.—*An alterative powder for a child 1 year old.*
- ℞ Potassii Iodidi, ℥ii; Decoctum Sarsæ Compositum ad ℥viii. M. Capiat ℥i ter in die.—*In rheumatism.*

℞ Tincturæ Guaiaci Ammoniatae, ℥iii ;
Tincturæ Aconiti, mxx ; Aquam Camphoræ ad ℥vi. M. ℥j ter in die.—*In tonsillitis and rheumatism.*

℞ Potassii Iodidi, ℥ss ; Spiritûs Ammoniæ Aromatici, ℥iii ; Tinct. Belladonnæ, ℥iss ;
Tincturæ Cinchonæ Compositæ, ℥vi ;
Aquam Menthæ Peperitæ ad ℥vi. M. ℥j ter in die.—*In some forms of asthma.*

℞ Potassii Iodidi, gr. xv ; Vini Colchici, ℥iss ;
Tincturæ Hyoscyami, ℥j ; Infusum Calumbæ ad ℥vi. M. Sumat ℥j ter in die.—*In chronic gout.*

℞ Potassii Iodidi, gr. xxvi ; Potassii Bromidi, ℥ss ; Ferri Ammoniæ Citratis, gr. xxx ;
Aquam Anethi ad ℥vi. M. ℥j ter in die.

℞ Hydrargyri Perchloridi, gr. i ; Potassii Iodidi, gr. lx ;
Tincturæ Calumbæ, ℥ii ; Aquam ad ℥vi. M. ℥j ter in die.—*In Constitutional syphilis.*

℞ Syrupi Ferri Iodidi, ℥iii ; Syrupi Aurantii, ℥ss ;
Aquam Anethi ad ℥vi. M. ℥j ter in die.

- R Potassii Bromidi, $\mathfrak{z}\text{ii}$; Ferri Ammoniae Citratis, $\mathfrak{z}\text{j}$; Spiritus Ammoniae Aromatici, $\mathfrak{z}\text{ii}$; Infusum Calumbæ ad $\mathfrak{z}\text{vi}$. M. $\mathfrak{z}\text{j}$ ter in die.—*In epilepsy.* (W. Fox.)
- R Magnesiae Sulphatis, $\mathfrak{z}\text{j}$; Tinct. Rhei, $\mathfrak{z}\text{ii}$; Syr. Zingiberis, $\mathfrak{z}\text{j}$; Aquam Carui ad $\mathfrak{z}\text{iss}$. M. $\mathfrak{z}\text{j}$ ter in die.—*An alterative for a child 1 year old.* (West.)
- R Liquoris Hydrargyri Perchloridi, $\mathfrak{z}\text{j}$; Liquoris Taraxaci, $\mathfrak{z}\text{ii}$; Extracti Cinchonæ Liquidi, $\mathfrak{z}\text{j}$; Tinct. Aurantii, $\mathfrak{z}\text{j}$; Syrupi, $\mathfrak{z}\text{iii}$; Aquam ad $\mathfrak{z}\text{iv}$. M. $\mathfrak{z}\text{ss}$ bis die.—*For a child 3 years old. An alterative and tonic in disorders of the liver.* (West.)
- R Extracti Taraxaci, $\mathfrak{z}\text{ii}$; Sodæ Bicarbonatis, $\mathfrak{z}\text{j}$; Extracti Sarsæ Liquidi, $\mathfrak{z}\text{iv}$; Syrupi Aurantii, $\mathfrak{z}\text{iv}$; Decoctum Sarsæ Ccpositum ad $\mathfrak{z}\text{vi}$. M. $\mathfrak{z}\text{ss}$ ter die in lacte.—*For a child 4 years old. Alterative and tonic in tabes mesenterica.* (West.)
- R Potassii Bromidi, gr. xii ; Tincturæ Belladonnæ, mxxv ; Syrupi, $\mathfrak{z}\text{ss}$; Aquam ad $\mathfrak{z}\text{iss}$. M. $\mathfrak{z}\text{i}$ ter in die.—*For a child 2 years old. In whooping-cough.*

℞ Hydrargyri cum Cretâ, gr. ii; Pulveris Ipecaeuanhæ Compositæ, gr. ss. M. One powder every six hours.—*In Inflammation of serous membranes. For an infant from 6—12 months.*

℞ Hydrargyri Iodidi Viridis, gr. xii; Extracti Lupuli, gr. lx; Ext. Opii, gr. v. M. Divide into xxiv pills. Sumat unam ter in die.—*In tubercular diseases of the skin, syphilis, &c.*

℞ Syrupi Ferri Phosphatis, ℥ii; Aquam ad ℥iss. M. ℥j ter in die.—*For a child a year old.*

℞ Ammonii Bromidi, gr. xii—lx; Infusi Aurantii, ℥vi. M. ℥j ter in die.—*In epilepsy. (Dr Gibb.)*

℞ Ammoniæ Benzoatis, gr. x—xxx; Aquæ ℥iss. M. Take three times a day.—*In inflammation of the bladder and phosphatic urine.*

ANTACIDS.

-
- R Sodæ Bicarbonatis, gr. x ; Spiritûs Ammoniaë Aromatici, ℥iii ; Syrupi, ℥ii : Aquam Anethi ad ℥iss. M. ℥j pro re nata.—*For a baby a week or more old.*
- R Bismuthi Carbonatis, ℥i ; Pulveris Aca-
ciæ, ℥iss ; Creasoti, ℥vi ; Spiritûs Am-
moniaë Aromatici ℥ii ; Syrupi ℥i ; Aquam
ad ℥viii. M. A sixth part three times
a day.—*In acidity of the stomach and
nausea.*
- R Lithiæ Carbon. gr. 5 ; Aquæ Anethi, ℥i.
M. Take three times a day.—*In chronic
gout. (Garrod.)*
- R Lithiæ Carbonatis, gr. v. Take in half
a bottle of soda water thrice daily.—
In gout.
- R Sodæ Bicarbonatis, gr. x ; Pulveris Rhei
compositæ, gr. xx ; Tincturæ Camphoræ
compositæ, ℥i ; Aquam Anethi ad ℥iss.
M.—*In acidity of the stomach with pain
and flatulence.*

- ℞ Liq. Bismuthi et Ammoniae Citratis, ℥i ;
Syr. Aurantii, ℥iii ; Infusum Chiratae
ad ℥iss. M.—*In pyrosis and gastro-*
dinia.
- ℞ Potassæ Bicarbonatis, ℥i ; Tincturæ
Hyoscyami, ℥vi ; Infusum Bucco ad ℥vi.
M. ℥i ter in die.—*In irritability of the*
bladder.
- ℞ Ammoniae Carbonatis, gr. v ; Potassæ
Chloratis, gr. x ; Tincturæ Opii, ℥x ;
Infusum Cinchonæ Flavæ ad ℥j. M.
Take three times a day.—*In debility*
with acidity.
- ℞ Tincturæ Lupuli, ℥vi ; Tincturæ Carda-
momi Co., ℥iv ; Vini Ipecacuanhæ, ℥ii ;
Extracti Opii Liquidum, ℥xxv ; Misturam
Cretæ ad ℥vi. M. ℥ss ter in die.—*In*
acidity and diarrhœa.
- ℞ Liquoris Potassæ, ℥xv ; Aquam ad ℥iss.
M. Ter in die.—*To render the urine*
alkaline.
- ℞ Potassæ Citratis, gr. xl ; Aquam ad ℥iii.
M. Take every three hours.—*Useful*
in the solvent treatment of uric acid
calculus. (Roberts.)

ANTISPASMODICS.

- R Tinct. Lobeliæ, ʒii ; Spt. Ætheris, ʒiii ; Tinct. Conii, ʒii ; Misturam Amygdalæ ad ʒvi . M. ʒss ter in die.—*In asthma.*
- R Folii Belladonnæ, gr. vi; Folii Hyoscyami, gr. iii; Folii Stramoni, gr. iii; Fol. Phellandrii Aquatici, gr. i; Ext. Opii, gr. $\frac{1}{4}$; Aquæ laurocerasi qs. Moisten the powdered leaves with the extract of opium dissolved in the laurel water, then dry and make into cigarettes.—*Most useful in asthma.*
- R Spiritûs Ætheris, ʒiii ; Spiritûs Chloroformi, ʒiii ; Tincturæ Cardamomi Compositæ, ʒvi ; Tincturæ Rheii, ʒiii ; Aquam Menthæ Viridis ad ʒvi . M. ʒj pro re nata.—*In flatulence.*
- R Potassæ Citratis, gr. exx; Tincturæ Stramonii, ʒi ; Tincturæ Colchici, ʒii ; Infusi Digitalis, ʒiii ; Aquam Menthæ Peperitæ ad ʒvi . M. ʒj ter in die,—*In gout, with dyspnœa and full pulse.*

- ℞ Tincturæ Valerianæ Ammoniatæ, ℥iv ; Spiritûs Chloroformi, ℥ii ; Infusum Valerianæ ad ℥vi. M. ℥j pro re nata.—*In hysteria.*
- ℞ Tincturæ Cardamomi Compositæ, ℥iv ; Acidi Hydrocyanici diluti, ℥xl ; Spiritûs Ammonia Aromatici, ℥ii ; Tincturæ Zingiberis, ℥iii ; Aquam Chloroformi ad ℥vi. M. ℥ss pro re nata.—*In simple colic.*
- ℞ Sodæ Carbonatis, gr. xii ; Spiritûs Ammonia Aromatici, ℥vi ; Syr. Simplicis ℥ii ; Tincturæ Cardamomi Compositæ, ℥xx ; Aquam Anethi ad ℥iss. M. ℥j quartâ-quâque horâ.—*In flatulence in babies 6 months old.*
- ℞ Zinci Sulphatis, gr. viii ; Extracti Belladonnæ, gr. ii ; Aquam ad ℥iv. M. ℥ss ter in die.—*In whooping-cough. Child over 3 years.*
- ℞ Tincturæ Assafœtidæ, ℥ii ; Tincturæ Castorei, ℥j ; Tincturæ Sumbulii, ℥ii ; Tincturæ Opii, ℥ss ; Aquam Menthae Peperitæ ad ℥viii. M. A sixth part every four hours.—*In hysteria.*

APERIENTS, CATHARTICS AND ANTHELMINTICS.

R Pulveris Rhei, \mathfrak{z} ss; Extracti Aloes Barbadensis, \mathfrak{z} ss; Ext. Belladonnæ, gr. iii; Olei Caryophylli, \mathfrak{m} iii. M. Divide into xx pills. One night and morning.
In habitual constipation.

R Extracti Aloes Barbadensis, gr. iii; Extracti Nucis Vomicae, gr. ss; Mastiche, gr. ss; Misce fiat pil. One pill night and morning.—*In habitual constipation in women. Very useful.*

R Magnesiae Sulphatis, \mathfrak{z} i—ii; Acidi Sulphurici Aromatici, \mathfrak{z} iss; Syrupi Zingiberis, \mathfrak{z} i; Aquam Chloriformi ad \mathfrak{z} vi. M. \mathfrak{z} j every two or three hours.—*A good aperient.*

R Magnesiae Sulphatis, \mathfrak{z} i—ii; Magnesiae Carbonatis, \mathfrak{z} ss; Sodæ Bicarbonatis, \mathfrak{z} j; Aquam Menthæ Viridis ad \mathfrak{z} vi. M. \mathfrak{z} j every two or three hours.—*A good aperient and antacid.*

℞ Magnesiae Sulphatis, ʒjss; Magnesiae Carbonatis, ʒss; Vini Colchici, ʒj; Aquam Chloroformi ad ʒvi. M. ʒj ter in die.—*In acute gout.*

℞ Calomelanos, gr. ii—gr. v; Pulveris Jalapae Compositae, gr. v—viii. M.—*Useful in thread worms in children.*

℞ Olei Crotonis, ʒi—ii; Olei Caryophylli, ʒi; Micæ panis, qs. M. Statim sumenda, et horis duabus repetenda, si opus sit.—*Useful where a brisk purge is required.*

℞ Olei Crotonis, ʒi vel ii; Pil. Coloeynthidis Co., ʒss; Pil. Assafætidae Co., ʒj. M. Divide in pilulas xviii. One or two at bedtime.—*In obstinate neuralgia.*

℞ Pulveris Rhei, gr. xv; Magnesiae Carbonatis, ʒj; Aquam Anethi ad ʒiss. M. ʒj every two or three hours in milk.—*An aperient for a child 12 months old.*

℞ Inf. Calumbæ ad ʒiv; Infusi Rhei, ʒivss; Tineturæ Aurantii, ʒii. M. ʒiii twice a day for a child three years old.—*Useful in threatened acute hydrocephalus.* (West.)

- ℞ Magnesiae Sulphatis, ℥ii; Syrupi Aurantii, ℥ii; Aquam Carui ad ℥iss. M. ℥ii every hour till the bowels act.—*For a child three years old.*
- ℞ Elaterii, gr. i; Extracti Gentianæ gr. xii. M. Divide into iv pills.—*One every night in dropsy.*
- ℞ Ext. Filicis Liquidii, ℥ss; Pulv. Acaciæ gr. xl; Spt. Myristicæ, ℥v; Syr. Tolutan ℥iv; Aquam Cinnamomi ad ℥j. M.—*In tape worm. For a child 8 or 10 years old, followed by a dose of castor oil.*
- ℞ Hydrargyri Subchloridi, gr. ii—iii; Pulv. Scammonii Co., gr. iv; Pulv. Cretæ Aromat, gr. v. M.—*At bed time for children with thread worm.*
- ℞ Santonini, gr. ii; Sacchari Lactis, gr. x. M.—*In round worm in children, given fasting, and followed by a brisk purge. For a child 3 years old.*
- ℞ Podophylli Resinæ, gr. i; Fellis Bovis, Pilulæ Hydrargyri, ā ā, gr. i; Pilulæ Rhei Compositæ, gr. ii. M. One pill at night.—*In obstinate constipation.*

- ℞ Kamelæ, gr. lx ; Syrupi, ℥ss ; Pulv. Trayaeanthæ, ℥j ; Aquam ad ℥iss. M. — *In tape worm; followed by a purge.*
- ℞ Podophylli Resinæ, gr. i ; Pilulæ Rhei Compositæ, gr. x ; Extracti Hyoseyami, gr. iv. M. Divide into iv pills. One every third night. — *In constipation.*
(Sir G. Burrows.)
- ℞ Pulveris Japalæ, ℥j ; Potassæ Tartratis Acidi, Ferri Carbonatis, ā ā, ℥j ; Pulv. Zingiberis, ℥ss. Theriacæ qs. ut fiat electuarium. Doses, ℥ss—℥j bis die. — *For worms in children.*
- ℞ Pulveris Rhei, gr. x ; Pulv. Scammonii, Potassæ Sulphatis, ā ā, gr. x ; Pulveris Cinnamomi Compositæ, gr. vi. Tere optime simul. Divide in pulveres vi. Sumat i. quartâquâque horâ ad alvi pluram solutionem. — *A mild aperient for young children.*
- ℞ Magnesiæ Sulphatis, gr. exx ; Mannæ, gr. elx ; Tincturæ Sennæ, ℥ij ; Infusum Sennæ ad ℥iss. M. Haustus. — *Common black draught.*

- R Pil. Cambogiæ Compositæ, gr. v ; Pil. Assafœtidæ Co., gr. v. M. Divide in pilulas ii. To be taken every night.—*A good purgative, acting chiefly on the small intestines.*
- R Pil. Colocynthis et Hyoscyami, gr. xx ; Fellis Bovini Purificati, Extracti Lupuli ā ā, gr. xx. M. Divide in pilulas xx. One three hours after dinner.—*In constipation and imperfect digestion.*
- R Sodæ Bicarb. ʒj ; Sodii Chloridi, gr. iv ; Sodæ Sulph. gr. x ; Mag. Sulph. gr. iij ; Ferri Sulphatis, gr. ʒ—l ; Aquæ Oj. M. —*In chronic gout. An imitation of Vichy water.*
- R Sodæ Tartaratae, ʒiv ; Tinct. Sennæ, ʒj ; Syrupi Zingiberis, ʒij ; Aquam ad ʒjss. M. Haustus.—*More pleasant than the common black draught.*
- R Confectionis Sulphuris, ʒij ; Extracti Taraxaci, ʒj. M. Dose.—*A tea-spoonful in the morning.*
- R Pilulæ Colocynthis Compositæ, gr. iiss ; Pilulæ Hydrargyri, gr. iiss. M.—*A good home aperient pill.*

- ℞ Elaterii, gr. iss; Pulveris Capsici, gr. ix; Hydrargyri Subchloridi, gr. xii; Ext. Hyoscyami, gr. xviii. M. Divide in pilulus xii.—*An active purgative. The capsicum prevents the nausea produced by elaterium.*
- ℞ Podophylli Resinæ, gr. iv; Pulveris Ipecacuanhæ, gr. vi; Extracti Gentianæ, gr. xxvi. M. Divide in pilulas xii.—*A useful pill in constipation.*
- ℞ Extracti Nucis Vomicæ, gr. iiij; Pulv. Ipecacuanhæ, gr. vi; Pilulæ Rhei Compositæ gr. xl. M. Divide in pilulas xii.—*An aperient pill in delicate women.*
- ℞ Pepsinæ Porci, gr. xxxii; Ext. Aloes Barbadensis, gr. viii; Glycerini, qs. M. Divide in pilulas viii.—*In some forms of dyspepsia with constipation.*
- ℞ Cusso, gr. 240; Mellis Depurati, ʒj. M. Take half at once and the rest in six hours.—*For tape worm.*
- ℞ Decocti Granati Radicis, ʒj—ij.—*In tape worm.*

ASTRINGENTS.

- R Olei Ricini, ʒj ; Pulveris Acaciæ, ʒss ; Sacchari Albi, ʒss ; Tincturæ Opii, ʒiv ; Aquæ Cinnamomi, ʒi. M. ʒj tertia vel quartâquâque horâ.—*In inflammatory diarrhœa in children. For a child 1 year old. (West.)*
- R Extracti Hæmatoxyli, ʒj ; Tincturæ Catechu, ʒii ; Syrupi, ʒj ; Aquæ Carui, ʒiss M. ʒi ter in die.—*In diarrhœa of children. For a child 1 year old. (West.)*
- R Liquoris Potassæ, ʒxxv ; Mucilaginis Acaciæ, ʒvi ; Vini Ipecacuanhæ, ʒxxiv ; Syrupi Tolutani, ʒiv ; Aquam ad ʒiv. M. ʒii sextaquâque horâ.—*In diarrhœa during teething. Child 12 to 18 months old.*
- R Acidi Sulphurici, ʒiss ; Tincturæ Opii, ʒj—iss ; Tincturæ Cinnamomi, ʒiii ; Syrupi, ʒs ; Aquam Camphoræ ad ʒvi. M. One ounce every time the bowels are moved.—*In severe diarrhœa, with vomiting. Early stages of cholera.*

- ℞ Tincturæ Catechu, ℥iii; Pulveris Cretæ Aromatici, ℥ss; Tincturæ Opii, ℥ss—i; Misturam Cretæ ad ℥vi. M. Sumat ℥i post singulas liquidas sedes.—*In simple diarrhæa.*
- ℞ Mag. Sulphatis, ℥i; Acidi Galliei, ℥i; Tinct. Digitalis, ℥j; Ext. Ergotæ, ℥ij; Infusum Rosæ Acidium ad ℥vi. M. ℥j every two or three hours.—*A valuable remedy in severe hæmoptysis.* (Dobell.)
- ℞ Tincturæ Matieo, ℥vi; Infusi Krameriæ, ℥viii; Syrupi Croci, ℥ii. M. ℥ss every three or four hours.—*In diarrhæa of phthisis.*
- ℞ Acidi Tannici, ℥ss; Acidi Nitrici dil. ℥j; Tincturæ Lupuli, ℥ii; Infusum Gentianæ ad ℥vi. M. ℥i ter in die.—*In chronic bronchitis, with profuse secretion.*
- ℞ Pilulæ Plumbi cum opio, gr. iv. To be taken every two or three hours.—*In passive hæmorrhages.*
- ℞ Tincturæ Opii, ℥j; Olei Cajuputi, ℥xii; Tincturæ Rhei, ℥ss; Misturam Cretæ Aromaticam ad ℥vi. M. ℥j post sedes liquidas.—*In simple painful diarrhæa.*

- ℞ Tincturæ Scillæ, ʒiss; Acidi Sulphurici diluti, ʒiss; Tinct. Opii, ʒss; Infusum Cascarillæ ad ʒvi. M. ʒss ter in die. — *To check profuse secretion in chronic bronchitis.*
- ℞ Chloroformi, ℥vi; Tincturæ Camphoræ Compositæ, ʒij; Olei Recini, ʒiii; Mucilaginis Tragacanthæ, ʒiii. M. — *To be taken at once in choleraic diarrhœa.*
- ℞ Alumenis Exsiccati, ʒj; Syrupi Rheados, ʒvi; Inf. Rosæ Acidum ad ʒviii. M. ʒj ter in die. — *In passive hemorrhage.*
- ℞ Acidi Gallici, gr. vi; Extracti Cannabis Indicæ, gr. $\frac{1}{2}$; Confectionis Rosæ, gr. i. M. Make into a pill, to be taken every night. — *In night sweats of phthisis.*
- ℞ Olei Terebinthinæ, ℥x—xx; Misturæ Amygdalæ, ʒj. M. To be taken every hour. — *In severe hæmoptysis in weak persons.*
- ℞ Olei Terebinthinæ, ʒij; Tincturæ Lavandulæ Compositæ, ʒss; Sacchari Purificati, ʒii; Pulveris Acaciæ, ʒii; Aquam ad ʒvi. M. ʒss statim et repetitur quartâquâque horâ si opus sit. — *In hemorrhage, typhoid fever, &c.*

- ℞ Extracti Belæ Liquidi, ʒij; Decocti Hordei, ʒiiss. M. Sumat post singulas sedes liquidas.—*In diarrhœa.*
- ℞ Cupri Sulphatis, gr. ss; Pulv. Opii, gr. ss; Confectionis Rosæ, qs. M. Fiat pil, ter die sumenda.—*In chronic diarrhœa and dysentery.*
- ℞ Pulveris Kino Compositæ, gr. x; Infusi Casearillæ, ʒvi; Aquæ Cinnamomi, ʒij; Syrupi Papaveris, ʒj. M. Haustus bis in dies sumendus.—*In chronic dysentery.*
- ℞ Misturæ Cretæ, ʒvi; Pulveris Cretæ Aromatiei, ʒj; Extracti Opii Liq. ℥xv; Aquam Cinnamomi ad ʒiv. M. A tea-spoonful every hour.—*In diarrhœa of children 1 year old.*
- ℞ Argenti Nitratis, gr. $\frac{1}{4}$ — $\frac{1}{2}$; Aquæ distillatæ, ʒij; Pulveris Acaciæ, gr. xl; Sacchari Albi, ʒij. M. ʒj bis vel tertîaquâque horâ.—*In obstinate diarrhœa.*
- ℞ Tineturæ Camphoræ Compositæ, ʒiii; Pulv. Cretæ Aromatiei, ʒii; Syrupi, ʒss; Aquam Camphoræ ad ʒvi. M. ʒj post sedes liquidas.—*In diarrhœa of typhoid.*

- ℞ Tinct. Ferri Perchloridi, ℥xv; Acidi Hydrochlorici Diluti, ℥x; Olei Menthæ Peperitæ, ℥j; Infusi Quassiae, ℥iss. M. Haustus ter in die.—*In hemorrhage from the stomach.*
- ℞ Zinci Oxidi, gr. xii; Ext. Conii, gr. xviii. M. Divide into six pills. One every night.—*For the relief of night sweats in phthisis.*
- ℞ Syrupi Papaveris, ℥iv; Bismuthi Subnitratis, ℥j; Mucil. Tragacanthæ, ℥iv; Aquam ad ℥viii. M. ℥iss ter in die.—*In the diarrhœa of phthisis.*
- ℞ Carbonis Ligni, ℥j; Theriacæ; ℥j. M. ℥j ter in die.—*In chronic diarrhœa with fecal fermentation.*
- ℞ Tinct. Ferri Pernitratis, ℥jss; Syrupi Simplicis, ℥ss; Aquam Anethi ad ℥vi. M. Sumat ℥j ter in die.—*In dysentery with the following pill.*
- ℞ Extracti Hæmatoxyli, gr. v; Spiritus Vini Rectificati, qs. Fiat pilula.
- ℞ Decocti Hæmatoxyli, ℥j; Acidi Nitrici Diluti, ℥x; Tincturæ Opii, ℥v. M. Haustus ter in die.

- ℞ Extracti Ergotæ Liquidi, ℥xl; Acid Gallici, gr. x; Aquæ Carui, ℥j. M. Haustus ter in die si opus sit.—*In severe hemorrhage from the stomach.*
- ℞ Bismuthi Subnitratis, gr. xv; Magnesiæ Carb., gr. xv; Mucilaginis Acaciæ, ℥ii; Aquæ Anethi, ℥j. M. Haustus.—*In dyspeptic diarrhœa.*

CAUSTICS AND COUNTER IRRITANTS.

-
- R Zinci Chloridi, $\bar{3}j$; Farina Tritici, qs.
Make a paste.—*Useful for cancer, &c.*
(Fell's Paste.)
- R Acidi Arseniosi, gr. xii; Hydrargyri
Subchloridi, $\bar{3}j$. M.—*In ulcerated lupus.*
Requires care. Dupuytren's powder.
- R Cupri Sulphatis, gr. x; Aquæ, $\bar{3}j$. M.
—*Useful over callous sores, syphilitic, &c.*
- R Hydrargyri, $\bar{3}j$; Acid Nitrici, $\bar{3}ii$. Solve.
For cancer, lupus, sloughing ulcers, &c.
Use a glass brush.
- R Iodinii, $\bar{3}j$; Potassii Iodidi, $\bar{3}ss$; Spiritûs
Vini Rectificati, $\bar{3}j$. M. Apply with
brush.
- R Antimonii Tartarati, gr. xl; Aquæ
Rosæ, $\bar{3}j$; Tincturæ Cantharidis, $\bar{3}j$. M.
—*Useful as a counter irritant to the chest*
in phthisis. Paint with a brush.
- R Zinci Chloridi, $\bar{3}j$; Zinci Oxidi, $\bar{3}j$. M.
Preserve in a well-stopped bottle.—
A caustic powder.

- ℞ Zinci Chloridi, gr. cccclxxx; Farinæ, gr. cxx vel qs; Liquoris Opii Sedativi, ʒj. M.—*Caustic paste.* (Lawson.)
- ℞ Argenti Nitratis, gr. x—xx; Aquæ Distillatæ, ʒj. M. To be applied with a brush to unhealthy sores.
- ℞ Antim. Chloridi, ʒj; Zinci Chloridi, ʒj; Farinæ Tritici, qs; Aquæ, qs. Make a paste.—*In lupus, &c.*
- ℞ Potassæ Causticæ, ʒj; Calcis, ʒj; Spt. Vini Rectificati, qs. Make a paste.—*In lupus, &c.*
- ℞ Argenti Nitratis, ʒj; Aquæ distil, ʒj. M.—*In rupia, impetigo, &c.*
- ℞ Arsenici Oxidi Albi, gr. x; Hydrargyri Sulphuratæ, gr. xl; Carbonis Animalis, gr. x. M. Côme's powder.—*In ulcerated lupus.*

DIAPHORETICS AND DIURETICS.

- ℞ Potassæ Nitratis, ʒj; Liq. Ammoniæ Acetatis, ʒss; Vini Ipecacuanhæ, ʒi; Sp. Ammoniæ Aromatici, ʒii; Decoctum Hordei ad ʒvi. M. ʒj ter in die.—*In catarrh.*
- ℞ Tincturæ Guaiaci Ammoniatæ, ʒiii-vi; Mucilaginis Tragacanthæ, ʒiii; Infusi Senegæ, ʒviii. M. ʒiss ter in die.—*Diaphoretic diuretic; stimulant and expectorant in chronic bronchitis.*
- ℞ Potassæ Nitratis, gr. cxx; Acidi Nitrici diluti, ʒi; Tinct. Digitalis, ʒss; Aquam Rosæ ad ʒvi. M. One ounce every four hours.—*In relapsing fever.*
(Murchison.)
- ℞ Ammoniæ Carbonatis, gr. v; Spiritûs Ætheris Chlorici, ℥x; Vini Colchici, ℥v; Liqueoris Ammoniæ Acetatis, ʒiii; Mucilaginis Acaciæ ad ʒiss. M. The draught to be taken every four hours.—*Useful in pneumonia complicated with gout.*

- R Potassæ Acetatis, \mathfrak{z} ss ; Aceti Scillæ, \mathfrak{z} ss ; Spiritûs Ætheris Nitrosi, \mathfrak{mxx} ; Tincturæ Digitalis, \mathfrak{mv} ; Decoctum Scoparii ad \mathfrak{z} iss. M. Ter die sumendus.—*In dropsy due to heart disease.*
- R Pulv. Opii, gr. j ; Potassæ Nitratis, gr. viii ; Pulveris Ipecacuanhæ Simplicis, gr. j. M. The powder at bedtime.—*In simple catarrh.*
- R Vini Ipecacuanhæ, \mathfrak{z} j ; Spiritûs Ætheris Nitrosi, \mathfrak{z} iss ; Potassæ Nitratis, \mathfrak{z} ss ; Syrupi Mori \mathfrak{z} ss ; Aquam ad \mathfrak{z} iss. M. \mathfrak{z} j ter in die.—*In a slight feverish attack. For a child from 1 to 2 years old.*
- R Vini Antimonialis, $\mathfrak{m}xlv$; Vini Ipecacuanhæ, \mathfrak{z} j ; Syrupi Rheados, \mathfrak{z} iii ; Liqueoris Ammonię Acetatis, \mathfrak{z} j ; Aquam ad \mathfrak{z} iii. M. \mathfrak{z} ii omni horâ.—*A depressing mixture for a child 2 or 3 years old.*
- R Vini Ipecacuanhæ, \mathfrak{z} iss ; Syrupi, \mathfrak{z} ss ; Tincturæ Camphoræ Compositæ, \mathfrak{z} ii ; Liqueoris Ammonię Citratis, \mathfrak{z} ss ; Aquam ad \mathfrak{z} ii. M. Sumat \mathfrak{z} j secundâ vel tertiâquâque horâ.—*In fever, catarrh, pneumonia, &c., in children from 3 to 5 years old.*

- R Tincturæ Scillæ, ʒij ; Tincturæ Camphoræ Compositæ, ʒvi ; Liquoris Ammoniac Citratis, ʒij ; Decoctum Scoparii ad ʒviii. M. Sumat ʒj ter in die.—*A diuretic for dropsy not due to kidney disease.*
- R Potassæ Acetatis, ʒiss ; Spiritûs Ætheris Nitrosi, ʒij ; Tinct. Digitalis, ʒj ; Tinct. Scillæ, ʒj ; Infusum Digitalis ad ʒvi. M. Sumat ʒss ter in die.—*In dropsy due to heart disease, and where there is diminished arterial pressure.*
- R Vini Antimonialis, ʒij ; Liq. Ammoniac Citratis, ʒiv ; Syrupi Hemidesmi, ʒiij ; Aquam ad ʒiv. M. Dose—one dessert-spoonful between 2 and 4 years.
- R Olei Copaibæ, ʒii ; Liq. Potassæ, ʒj ; Tincturæ Lavandulæ Compositæ, ʒiij ; Aquam Camphoræ ad ʒviii. M. ʒj ter die.—*In gonorrhœa.*
- R Pulveris Digitalis ; Pulveris Scillæ, ā ā gr. xii ; Extracti Taraxaci, gr. xxxvi. M. Divide into xii pills ; one to be taken twice daily.—*As a diuretic in mitral, but dangerous in aortic disease. (Dr Tanner.)*

- ℞ Pilocarpin Hydrochloratis, gr. iv; Aquæ
 ʒj. M. Five minims subdermically.—
*Used with great success in the case of a
 child 3 years old; œdema of the lung.*
 (A. Weber.)
- ℞ Olei Juniperi, ℥xx; Syr. Hemidesmi, ʒss;
 Mucilaginis Acaciæ, ʒiv; Decoctum
 Taraxaci ad ʒviii. M. ʒi ter in die.—
Diuretic and diaphoretic.
- ℞ Acidi Sallicylici, ʒj; Liquoris Ammoniac
 Acetatis, ʒii; Aquam Chloroformi
 ad ʒvi. M. ʒj quartâquâque horâ.—
In rheumatic fever.
- ℞ Sodæ Sallicylici, ʒiss; Syr., ʒj; Aquam
 ad ʒvi. M. Two table-spoonfuls every
 two hours for four doses, and then
 every four hours.—*In rheumatic fever.*
- ℞ Liq. Taraxaci, ʒiij; Syr. Aurantii, ʒiv;
 Potassii Iodidi, gr. xii; Spt. Etheris
 Nitrosi, ʒj; Potassæ Nitratis, gr. xxx;
 Tinct. Scillæ, ℥xxx; Tinct. Digitalis,
 ℥xxiv; Aquam ad ʒviiij. M. ʒss
 quartâquâque horâ.—*A good diuretic
 for a child 6 years old.*

EMETICS AND EXPECTORANTS.

- ℞ Vini Ipecacuanhæ, ʒj. To be given at once and followed by a tea-cupful of warm water.—*An emetic for children.*
- ℞ Cupri Sulphatis, gr. x; Aquæ, ʒiii. M. To be given at once, and repeated, if necessary, in half an hour.
- ℞ Ammonię Carbonatis, gr. xii; Tinct. Scillæ, ℥xvi; Syr. Tolut, ʒiii; Infusum Senegę ad ʒiii. M. ʒii quartâquâque horâ. *In chronic bronchitis. For a child 2 or 3 years old.*
- ℞ Vini Ipecac., ʒj; Spt. Ætheris Nitrosi, ʒj; Liq. Ammonię Acetatis, ʒj; Syr., ʒii; Aquam Anethi ad ʒiss. M. ʒj quartâquâque horâ.—*Early stages of bronchitis. For a child 1 year old.*
- ℞ Zinci Sulphatis, gr. xx—xl; Aquam ad ʒiii. M.—*An emetic in poisoning.*
- ℞ Pulveris Sinapis, ʒss; Aquam ad ʒiii. M.—*Most useful emetic in alcoholic poisoning.*

- ℞ Ammoniae Carbonatis, ʒss; Spiritûs Ætheris, ʒiii; Tincturæ Scillæ, ʒiss; Tincturæ Camphoræ Compositæ, ʒiii; Tincturæ Lavandulæ Compositæ, ʒiv; Infusum Senegæ ad ʒviii. M. ʒj ter in die.—*In chronic bronchitis.*
- ℞ Succī Conii, ʒj; Acidi Hydrocyanici diluti, ℥x; Vini Ipecacuanhæ, ʒj; Syrupi ʒss; Aquam ad ʒiss. M. Sumat ʒj quartâquâque horâ.—*In whooping-cough. For a child 2 years old.*
- ℞ Tincturæ Lobeliæ Æthericæ, ʒiii; Syrupi Rheados, ʒvi; Succī Conii, ʒii; Misturam Amygdalæ ad ʒvi. M. ʒj every four hours.—*In spasmodic cough and asthma.*
- ℞ Vini Ipecacuanhæ, ℥xx; Spiritûs Ammoniae Aromatici, ℥v; Syrupi, ʒiii; Aquam Anethi ad ʒiss. M. ʒj ter in die.—*For a child 6 months old.*
- ℞ Pilulæ Scillæ Compositæ, Extracti Conii ā ā ʒss. M. Divide into 12 pills. One or two daily.—*In chronic cough.*
- ℞ Infusi Senegæ, ʒiii; Misturæ Amygdalæ, ʒvi; Tincturæ Opii, ℥viii. M. ʒj ter in die.—*Chronic cough.*

- R Vini Ipecacuanhæ, ʒii; Aquæ ʒj; Mucilaginis Acaciæ, ʒss; Syrupi Limonis, Syrupi Papaveris ā ā ʒj. M. ʒi—ʒii secundis horis.—*For catarrh in children.*
- R Infusi Senegæ, ʒj; Liquoris Ammoniac Acetatis, ʒss; Syrupi Scillæ, ʒii; Syrupi Papaveris, ʒii. M. ʒj ter in die.—*In bronchitis in young children.*
- R Tincturæ Camphoræ Compositæ ʒiii; Vini Ipecacuanhæ, ʒii; Liquoris Ammoniac Acetatis, ʒss; Infusum Gentianæ ad ʒvi. M. Sumat ʒj ter in die.—*Expectorant and tonic.*
- R Pilulæ Scillæ Compositæ, gr. v. Unam nocte manequē.—*Expectorant in chronic bronchitis.*
- R Oxymelis Scillæ, ʒii; Syr. Papaveris, ʒi; Tincturæ Camphoræ Compositæ, ʒss; Tinet. Ætheris Lobeliæ, ʒss; Misturæ Ammoniacy, ʒj. M. Haustus.—*In the fit of asthma.*
- R Oxymelis Scillæ, ʒj; Vini Antim. ʒii. M. A tea-spoonful every 10 or 15 minutes till sickness is produced.—*In catarrhal affections of children.*

- R Ammonii Chloridi, gr. x; Spt. Chloroformi, ℥xx; Vini Antimonialis, ℥x; Aquam Camphoræ ad ℥iss. M. Fiat Haustus. Sumat ter in die.
- R Tincturæ Camphoræ Compositæ, ℥ss; Syrupi Tolutani, ℥j; Misturam Ammoniaci ad ℥vi. M. ℥ss ter in die.
- R Ammoniae Bicarbonatis, ℥j; Tincturæ Digitalis, ℥j; Tinct. Camphoræ Compositæ, ℥iij; Mist. Amygdalæ ad ℥vi. M. Sumat ℥j ter in die.—*In chronic bronchitis, with heart disease.*
- R Tincturæ Camphoræ Compositæ, ℥iij; Spiritûs Ætheris, ℥ij; Tinct. Scillæ, ℥j; Syrupi, ℥ss; Aquam ad ℥vi. M. ℥j ter in die.—*Bronchitis and asthma.*
- R Succī Belladonnæ, ℥j; Acidi Hydrocyanici diluti, ℥x; Viui Ipecac., ℥j; Syrupi, ℥ss; Aquam ad ℥iss. M. ℥j ter in die.—*In whooping-cough. Child from 2 to 3 years old.*
- R Ammonii Bromidi, gr. xii; Vini Ipecacuanha, ℥j; Succī Conii, ℥ss; Syrupi, ℥iij; Aquam ad ℥iss. M. ℥j ter in die.—*In whooping-cough or other spasmodic cough. Child from 1 to 2 years.*

GARGLES AND INHALATIONS.

- R Alumenis exsiccati, ʒj ; Infusum Rosæ Acidum ad ʒvi. M.—*A simple gargle in relaxed throat.*
- R Potassæ Chloratis, ʒiss ; Acidi Hydrochlorici, ℥xl ; Glycerini, ʒj ; Aquam ad ʒxx. —*A gargle in scarlet fever.*
- R Liq. Sodæ Chloratæ, ʒiv ; Aquam ad ʒviii. M.—*A gargle in scarlet fever.*
- R Zinci Sulph., gr. xv ; Syrupi Mori, ʒiv ; Glycerini, ʒj ; Inf. Krameriæ ad ʒviii. M.—*In relaxed throat.*
- R Alumenis, ʒii ; Tincturæ Myrrhæ, ʒj ; Aquam ad ʒviii. M.—*In mercurial salivation.*
- R Alumenis, ʒiss ; Tinct. Capsici, ʒii ; Inf. Acidi Rosæ ad ʒviii. M.—*In relaxed uvula and tonsils.*
- R Creasoti ℥xx ; Tinct. Lavand. Co., ʒiv ; Tinct. Myrrhæ, ʒiv ; Syrupi, ʒiss ; Aquam ad ʒviii. M.—*In clergyman's sore throat.*

- ℞ Acidi Tannici, ℥ss ; Glycerini, ℥iv. M.
Paint the throat with a brush.—*In relaxed throat when gargling is difficult.*
- ℞ Sodæ Biboratis, ℥j ; Glyccrini, ℥ii ;
Aquæ Rosæ, ℥iv. M.—*Paint over mouth in aphthæ.*
- ℞ Sodæ Sulphitis, gr. lx ; Aquæ, ℥j. M.—
In aphthæ.
- ℞ Hydrargyri Perchloridi, gr. ii ; Acidi
Nit. dil, ℥xxx ; Tinct. Myrrhæ, ℥j ;
Aquam ad ℥viii. M.—*In syphilitic affec-
tions of the mouth.*
- ℞ Vapor Conii, ℥j B.P. Take xx drops,
place in a vessel of boiling water and
inhale the steam.
- ℞ Creasoti, ℥xxx ; Aquæ Bullicntis Oss.
—*In phthisis, ozæna, &c.*
- ℞ Olei Terebinthinæ, ℥j ; Aquam Calidam
ad ℥vi. M.—*As an inhalation in chronic
bronchitis with great secretion.*
- ℞ Acidi Hydrocyanici dil, ℥xx ; Tinct.
Hyoscyami, ℥j ; Tinct. Lupuli, ℥j ;
Aquam Calidam ad ℥viii. M.—*As an
inhalation in phthisis, ulceration of
larynx, &c.*

LOTIONS, LINIMENTS & OINTMENTS.

-
- ℞ Spiritûs Rectificati, ʒj ; Acidi Acetici diluti, ʒj ; Ammoniaë Hydrochloratis, ʒiii ; Aquæ Oj. M.— *A good lotion for orchitis, &c.*
- ℞ Plumbi Nitratis, ʒj ; Aquam ad. Oj. M. *A good lotion for foul ulcers, cancers, &c.*
- ℞ Extracti conii, ʒj ; Aquam ad ʒiii. M. *Macerate for two hours, and strain.— For painful sores.*
- ℞ Liquoris Zinci Chloridi, ʒii ; Aquam ad ʒviii. M.— *The best lotion for cancer of the womb.*
- ℞ Acidi Carbolici, ʒj ; Glycerini, ʒj ; Aquam ad ʒviii. M.— *For sores, or injecting the womb after bad delivery.*
- ℞ Vini Colchici, ʒss ; Tincturæ Opii, ʒss ; Sodæ Carbonatis ʒss. M.— *To be sprinkled on clothes wrung out of hot water, and applied to the joints in gout.*

- ℞ Boracis, gr. lx—cxx ; Glycerini, ℥j ;
Aquam Sambuci ad ℥viii. M.—*Lotion
for squamous diseases of the skin.*
- ℞ Alumenis, gr. iv ; Aquam ad ℥j. M.—*A
useful lotion for inflammation of the eyes.
Use two or three drops twice a day, or
oftener if necessary.*
- ℞ Acidi Hydrocyanici diluti, ℥iii ; Plumbi
Acetatis, gr. lx ; Spiritus Rectificati, ℥i ;
Aquam Sambuci ad ℥viii. M.—*Useful
in prurigo, &c., with much itching. The
skin must not be wounded.*
- ℞ Liquoris Ammoniae Acetatis, ℥j ; Acidi
Hydrocyanici dil., ℥iss ; Enemæ Tabaci
ad ℥viii. M.—*Sponge the part twice a
day. Useful in pruritus of the anus and
vulva.*
- ℞ Liq. Plumbi Subacet., ℥j ; Glycerini, ℥ss ;
Aquam ad ℥vi. M.—*As a lotion in
acute ekzema.*
- ℞ Tinct. Cantharidis, ℥j ; Glycerini, ℥iss ;
Acidi Acetici, ℥iss ; Spiritus Rose-
marini, ℥iss ; Aquæ Rosæ, ℥viii. M.—
*A most valuable application in alopecia
not of parasitic origin.*

- ℞ Sodæ Hyposulphitis, ʒii ; Glycerini, ʒss ; Aquam Rosæ ad ʒiv. M.—*A lotion in tinea.*
- ℞ Liq. Carbonis Detergentis, ʒss ; Acidi Hydrocyanici diluti, ʒj ; Glycerini, ʒii ; Aquam ad Oss. M.—*Useful in ekzema and pruritus.*
- ℞ Acidi Sulphurosi, ʒiv ; Aquam distillatam ad ʒviii. M.—*A lotion in all parasitic diseases.*
- ℞ Zinci Sulph., gr. xxx ; Zinci Oxidi, gr. lx ; Pulveris Acaciæ, ʒii ; Aquam Sambuci ad Oj. M.—*Useful in ekzema.*
- ℞ Atropiæ Sulphatis, gr. ii ; Aquæ Distillatæ, ʒss. M.—*A drop to dilate the pupil in iritis.*
- ℞ Linimenti Camphoræ Compositæ ʒiss ; Linimenti Belladonna, ʒiss. M.—*Useful in lumbago.*
- ℞ Lin. Belladonnæ ; Lin. Aconiti ā ā, ʒiss. M.—*Useful in neuralgia, pleurodynia, &c.*
- ℞ Linimenti Belladonnæ ; Lin. Opii ā ā, ʒj ; Lin. Terebinthinæ, ʒii. M.—*Chronic rheumatism, &c.*

- ℞ Linimenti Opii, ʒj; Linimenti Bella
donnae; Linimenti Hydrargyri ā ā, ʒi.
M.—*In syphilitic tubercles, nodes, &c.*
- ℞ Olei Morrhuæ, ʒiiiss; Spiritus Am-
moniae Aromatiei, ʒi; Tinet. Opii, ʒiv;
Olei Cajuputi, ʒi. M.—*To be rubbed
over the abdomen when cod liver oil
cannot be borne by the stomach.*
- ℞ Bismuthi Subnitratis, ʒj; Unguenti
Simplieis, ʒi. M.—*A useful ointment in
strumous sores.*
- ℞ Aeidi Carboliei, gr. lx; Unguenti Zinci
ʒi. M.—*Useful in ekzema, psoriasis, &c.*
- ℞ Camphoræ, gr. lx; Spiritus vini Reeti-
ficati, qs; Calomelanos, gr. xl; Adepis
puræ, ʒi. M.—*In pruritus of vulva, &c.*
- ℞ Plumbi Acetatis, gr. x; Zinei Oxidi,
gr. xx; Hydrargyri Subehloridi, gr. xx;
Unguenti Hydrargyri Nitratis, gr. xx;
Adipis Recentis, ʒss; Olei Palmæ
Purificati, ʒss. M.—*In ekzema capitis,
&c.*
- ℞ Unguenti Sulphuris, ʒj; Tinet. Lavan-
dulæ Compositæ, ʒj. M.—*In scabies.*

- R Acidi Hydrocyanici Diluti, $\text{m} \times \text{l}$; Olei Cadini, ̄j ; Saponis Mollis, ̄ij ; Olei Rosmarini, ̄jss ; Aquam ad ̄viii . M. Fiat Lotion. To be rubbed over the parts affected night and morning.—*Chronic ekzema, psoriasis.*
- R Olei Amygdalæ dulcis, ̄j ; Liquoris Ammoniae Fortioris, ̄j ; Spiritus Rosmarini, ̄iv ; Aquæ, ̄ij . M.—*A stimulant lotion for the hair.*
- R Atropiæ Sulphatis, gr. j; Boracis, gr. 120; Glycerini, ̄ss ; Acidi Hydrocyanici Diluti, ̄j ; Aquæ Floris Aurantii, ̄ij ; Aquam ad ̄xii . M.—*Useful to allay itching, &c.*
- R Calaminæ, ̄j ; Cretæ Præparatæ, ̄j ; Glycerini, ̄iij ; Acidi Hydrocyanici dil., ̄ss ; Liquoris Calcis, ̄iij ; Aquam Rosæ ad ̄vii . M. Fiat Lotion.—*In ekzema and acne. (Startin.)*
- R Olei Succini, ̄ss ; Olei Olivæ, ̄ss ; Olei Caryophylli, ̄j . M. Fiat Lini-
mentum.—*A substitute for Roche's embrocation. Useful in whooping-cough, when rubbed down the spine.*

- ℞ Creasoti, ℥vi; Ung. Hydrargyri, gr. xxx; Hydrargyri Oxidi rubri, gr. xx; Adipis Recentis, ℥j. M.—*In parasitic and other skin diseases.*
- ℞ Cretæ Preparatæ, ℥j; Adipis Benzoati, ℥j. M.—*In ekzema, when more stimulating applications irritate.*
- ℞ Glycerini, Acidi Carbolici ā ā ℥ij; Sulphuris Præcipitati, Spiritûs, Vini Rectificati ā ā ℥ij. M.—*To be applied every night in acie rosacea. (Hebra.)*
- ℞ Ungenti Zinci Benzoati, ℥j; Spiritûs Vini Rectificati, ℥ij. M.—*In ekzema. (Erasmus Wilson.)*
- ℞ Boracis, ℥j; Liquoris Plumbi Subacitatis, ℥xxx; Glycerini, ℥ij; Aquam ad ℥viii. M. Fiat Lotion.—*In ekzema of the face and eyelids.*
- ℞ Aluminis, gr. xxiv; Atropiæ Sulp. gr. j; Aquæ ℥viiij. M. Fiat Lotion.—*In iritis.*
- ℞ Argenti Nitratis, gr. i—iv; Aquæ distillatæ, ℥j. M.—*In conjunctivitis, &c.*

NARCOTICS AND SEDATIVES.

- R Liquoris Morphiae, ℥iiss; Acidi Phosphori diluti, ℥i; Acidi Hydrocyanici Diluti, ℥xii; Syrupi Rheados, ℥ss; Aquam ad ℥iiss M. ℥j tusseurgenti.—*A most useful cough medicine in phthisis.*
- R Chloral Hydratis, gr. x; Potassii Bromidi, gr. x; Syrupi Aurantii, ℥ss; Aquæ, ℥j. M. To be taken at bedtime.—*A good sedative in nervous excitement. May be given thrice daily.*
- R Extracti Cannabis Indicæ, gr. $\frac{1}{4}$ —gr. i; Confectionis Rosæ, qs. M. Fiat pilula. Take one at bedtime. *In sleeplessness of neuralgic headache. (Day.)*
- R Tinct. Opii, ℥j; Potassii Bromidi, ℥iiss; Syrupi Rheados, ℥j; Aquam ad ℥vi. M. Sumat ℥j ter in die.
- R Tinct. Opii, ℥j; Mucilaginis Acaciæ, ℥ii; Syr. Simp., ℥ii; Aquam Anethi ad ℥j. M. ℥i bis terve in die.—*For infants.*

- ℞ Tincturæ Belladonnæ, ʒj; Tincturæ Conii, ʒii; Vini Ipecacuanhæ, ʒiss; Aquam Chloroformi ad ʒvi. M. ʒj ter in die. *A sedative in irritable coughs.*
- ℞ Tincturæ Camphoræ Compositæ, ʒiij; Vini Ipecacuanhæ, ʒj; Acidi Hydrocyanici diluti, ℥xii; Syrupi Scillæ, ʒss; Aquam ad ʒvi. M. ʒj ter in die.—*A good cough mixture.*
- ℞ Pulveris Ipecacuanhæ Compositæ, gr. i; Sacchari albi, gr. xx. M. Divide into four powders. One to be given occasionally.—*For infants from 1 to 4 weeks old.*
- ℞ Tincturæ Digitalis, ℥x—xv; Acidi Hydrocyanici diluti, ℥iii; Tincturæ Opii, ℥v; Aquam Camphoræ ad ʒj. M. Take three times a day.—*Sedative in heart disease.*
- ℞ Liquoris Morphię Hydrochloratis, ℥xx; Chloroformi, ℥x; Tincturæ Cannabis Indicæ, ℥xx; Pulveris Tragacanthæ Compositæ, gr. xxx; Spt. Ætheris, ℥xl; Acidi Hydrocyanici diluti, ℥iv; Aquam ad ʒiss M. Haustus. *A substitute for chlorodyne.*

- R Tincturæ Opii, $\bar{5}$ iss; Spiritûs Etheris Nitrosi, $\bar{5}$ ii; Aquam Camphoræ ad $\bar{3}$ vi. M. Sumat $\bar{3}$ j quartâquâque horâ.—*To check after-pains.*
- R Liquoris Morphiae Hydrochloratis \mathfrak{m} xv; Spiritûs Ammoniae Fœtidi, $\bar{5}$ ss; Aquam Camphoræ ad $\bar{3}$ iss. M. Hanstus statim sumendus.—*To prevent nervous shock after labour.*
- R Extracti Hyoscyami, Camphoræ, Lupulinæ \bar{a} \bar{a} gr. xx. M. Divide into xii pills. Two at bedtime.—*In sleeplessness in hysterical patients.*
- R Codeiæ, gr. $\frac{1}{2}$; Pilulæ Assafœtidæ Compositæ, gr. v. M. Fiat pilula.—*At bedtime in spasmodic cough. (Tanner)*
- R Tinct. Veratri Viridis, \mathfrak{m} v; Aquæ, $\bar{3}$ j. M. To be taken every three hours, increasing the tincture by a drop till the pulse is sufficiently lowered.—*In pneumonia, pleurisy, acute rheumatism.*
- R Extracti Opii, gr. i vel iii; Extracti Belladonnæ, gr. $\frac{1}{3}$; Butyri Cacao, gr. xx. M. Make into a suppository.—*In diseases of the bladder and rectum.*

R Calcis Hypophosphatis, ʒj; Tincturæ Camphoræ Compositæ ʒiij; Acidi Hydrocyanici, ʒxii; Syrupi Simplicis, ʒj; Infusum Calumbæ ad ʒvi. M. ʒj ter in die. *Very useful in phthisis.*

R Ext. Opii, gr. i—ii; Pulv. Capsici, gr. ii; Extracti Hyoseyami, gr. iv. M. Divide in pilulas duo. Take one at bedtime. —*Most useful in vomiting, and where opium is required.* (Sir James Simpson.)

R Tincturæ Camphoræ Compositæ, ʒxx; Vini Ipecacuanhæ, ʒx; Vini Antimonialis, ʒxxx; Mist. Amygdalæ ad ʒj. M. ʒii quartâquâque horâ. —*For a child 1 year old.*

R Vini Ipecac., ʒx; Oxymel Scillæ, ʒxl; Spiritûs Etheris Nitrosi, ʒxx; Tinct. Camphoræ Compositæ, ʒxx; Aquæ Anisi ʒiss. M. ʒii ter in die. —*Cough mixture for children 1 year old.*

REFRIGERANTS AND SALINES.

- R Potassæ Bicarbonatis, gr. xl; Acidi Citrici, gr. xx; Vini Antimonialis, ℥iss; Vini Ipecac., ℥xx; Syr. Limonis, ℥iiss; Aquam ad ℥iii. M. ℥ii every three or four hours.—*For a child 2 years old. In the early stages of croup after the child has been placed in a hot bath. (West.)*
- R Quiniæ Sulphatis, gr. xii; Pulveris Digitalis, gr. xii; Pulveris Opii, gr. vi. M. Divide into twelve powders and give one every four hours.—*Niemeyers powders, in acute phthisis with high temperature.*
- R Liquoris Ammoniaë, ℥ss; Vin Ipecacuanha, ℥j; Potassæ Nitratis, ℥ss; Syr. ℥j; Sacchari Usti, ℥ss; Aquam Camphoræ ad ℥vi. M. ℥j ter in die.—*In acute catarrh.*
- R Potassæ Bitart. ℥j; Olei Limonis, ℥xv; Sacchari Albi, ℥ii; Aquam Bullientem ad ℥xl. M.—*A pleasant drink when the thirst is great in fevers.*

- ℞ Potassæ Chloratis, ʒj; Syr. Limonis, ʒiii;
Aquam ad Oj. M.—*As a drink in scarlet
fever. Half to a wine-glassful for a
child.*
- ℞ Potassæ Nitratis, gr. xii; Magnesiae
Sulphatis, ʒj; Syrupi Limonis, ʒiii;
Aquam ad ʒiss. M. ʒii ter in die.—
*Febrifuge and aperient. For a child a
year old.*
- ℞ Potassæ Bicarbonatis, gr. xx; Aeidi
Citrici gr. xx; Vini Ipecacuanhæ ℥xi;
Tincturæ Hyoscyami ℥xviii; Syrupi
Mori, ʒiii; Aquam ad ʒiss. M. ʒii every
six hours.—*For a child a year old. A
useful saline and sedative for children.*
- ℞ Potassæ Bicarb. ʒii; Spiritus Ammoniae
Aromatisei, ʒij; Syrupi Aurantii, ʒj;
Aquam ad ʒvi. M.
Aeidi Citrici, ʒiss; Aquam ad ʒvi. M.
Two table-spoonfuls of each to be mixed
together and drank during effervescence
every four hours.
- ℞ Acidi Citrici, Potassæ Bicarb. ā ā, ʒj;
Syr. Aurantii, ʒj; Spt. Myristicæ, ʒj;
Aquam ad ʒvi. M. ʒj ter in die.—
Inflammatory sore throat.

- ℞ Acidi Hydrochlorici dil, ʒiii; Mellis Depurati, ʒj; Dococt Hordei Oii. M.—*As a daily drink in typhus, &c.*
- ℞ Liquoris Ammonia, Acetatis ʒvi; Spiritus Ammonia Aromatici, ʒiii; Syrupi Aurantii, ʒj; Aquam ad ʒvi. M. ʒj ter in die.—*In early stages of fever, catarrh, &c.*
- ℞ Potassæ Chloratis, gr. xl; Acidi Hydrochlorici, ʒj; Spiritus Chloriformi, Liq. Cinchonæ ā ā, ʒj; Aquam ad ʒvi. M. ʒss ter in die.—*In herpes zoster.*
- ℞ Potassæ Nitratis, gr. xx; Ammonia Chloridi, gr. xii; Aquæ, ʒiss. M. Bis, ter, quaterve in die.—*In inflammatory fever.*
- ℞ Vini Antimonialis, ʒss; Liq. Ammonia Acetatis, ʒss; Mag. Sulphatis, ʒj; Aquam Camphoræ ad ʒvi. M. Sumat ʒj ter in die.—*Diaphoretic and aperient.*
- ℞ Potassæ Nitratis, gr. xl; Sodæ Bicarbonatis, ʒss; Vini Antimonialis, ʒii; Syrupi Croci, ʒj; Spiritus Aetheris Nitrosi, ʒj; Aquæ, ʒv. M. ʒiss ter die. *A good febrifuge.*

- ℞ Potassæ Nitratis, ʒj; Spiritûs Ætheris Nitrosi, ʒiv; Syrupi, ʒss; Decocti Taraxaci ad ʒvi. M. ʒj ter in die.
- ℞ Antimonii Taratarati, gr. i; Potassiæ Nitratis, ʒii; Tinct. Hyoscyami, ʒiss; Misturæ Amygdalæ, ʒxii; Tincturæ Digitalis, ʒss. M. ʒj every hour.—*In bronchitis and dropsy of the chest.*
- ℞ Liquoris Ammonię Acetatis, ʒss; Spiritûs Ammonię Aromatici, ʒvi; Vini Ipecacuanhæ, ʒss; Syrupi, ʒiii; Aquam ad ʒiss. M. ʒj ter in die.—*For a child 6—12 months old. In slight fever.*
- ℞ Liquoris Ammonię, ʒj; Ammonię Carbonatis, ʒss; Potassæ Nitratis, ʒj; Aquam Camphoræ ad ʒvi. M. Sumat ʒj ter in die.—*In simple catarrh.*
- ℞ Succı Limonis, ʒiv; Cort. Limonis recentis conc., ʒss; Potass. Nitrat., ʒj; Aquæ ferventis, Oij. M. Stent ad refrigerationem et cola.—*A common drink in fevers.*

STIMULANTS.

- ℞ Ammoniae Bicarbonatis, ℥ss ; Spiritus Myristicae, ℥ij ; Syrupi, ℥j ; Infusum Gentianae Compositum ad ℥vi. M. ℥j quartâquâque horâ.
- ℞ Spt. Ætheris, ℥iss ; Spt. Ammoniae Aromatici, ℥ij ; Infusum Cinchonae ad ℥vi. M. ℥j ter in die.
- ℞ Misturæ Spiritus Vini Gallici, ℥vi. M. ℥ss—℥j pro re nata.—*A valuable stimulant.*
- ℞ Vini, ℥vi ; ovor duor vetellos ; Sacchari Purificati, ℥ss ; Olei Cinnamomi, ℥iv ; Tincturæ Capsici., ℥xx. M. ℥iss ter, quaterve aut sæpius in die.—*In debility without fever.*
- ℞ Liqueur Sodæ Chloratæ, ℥j—ii ; Syrupi Tolutani, ℥j ; Tinct. Serpentariæ, ℥vi ; Aquam ad ℥viii. M. A sixth part every six hours.—*In low fever, with furred tongue and offensive evacuations. (Dr Tanner.)*

- ℞ Vini Xerici, ℥ii; ovi vetellum unius;
Jusculi Bovini (sine sale), ℥viii. M. Sit
enema.—*A restorative in fevers.*
- ℞ Spiritûs Chloroformi, ℥vi; Misturæ
Spiritûs Vini Gallici, ℥viii. M. A sixth
part every four hours.—*Low stages of
fever.*
- ℞ Ovor. Vetellos duor; Spiritûs Vini
Galliei, ℥j; Sac. Albi, qs; Lactis, ℥viii.
M.—*In inertia of the uterus during
labour.*

TONICS.

- ℞ Misturæ Ferri Compositæ, ℥iv ; Tincturæ Scillæ, ℥xvi ; Tinct. Conii, ℥xl ; Misturam Amygdalæ ad ℥iii. M. ℥ii ter in die.—*In debility during convalescence from whooping-cough ; whoop still present. Child 2 years old. (West.)*
- ℞ Acidi Hydrocyanici diluti, ℥viii ; Extracti Cinchonæ Liquidi, ℥iss ; Syrupi Aurant. ℥iss ; Aquæ Flor. Aurant. ℥iii ; Aquam Distil ad ℥iss. M. ℥ii ter in die.—*In whooping-cough when iron disagrees. Child 2 years of age. (West.)*
- ℞ Ammoniæ Carbonatis, gr. xxx ; Tincturæ Aconiti, ℥xl ; Tinct. Cinchonæ, ℥vi ; Aquam Menthæ Piperitæ ad ℥vi. M. ℥j ter in die.—*In pneumonia.*
- ℞ Acidi Nitrici diluti, ℥xvi ; Acidi Hydrochlorici diluti, ℥xxiv ; Acidi Hydrocyanici diluti, ℥viii ; Spiritûs Chloroformi, ℥xl ; Tincturæ Aurantii, ℥iss ; Syr. Simplicis, ℥ii ; Aquam ad ℥iv. M. ℥ss ter in die.—*In phthisis, with cod liver oil. Child 4 years old.*

- ℞ Quiniæ Sulphatis, gr. xii; Acidi Phosphorici diluti, ℥iss; Tinct. Lupuli, ℥vi; Aquam ad ℥viii M. ℥j ter in die.
- ℞ Sodæ Bicarbonatis, gr. xxiv; Tincturæ Rhei, ℥j; Extracti Taraxaci, gr. xl; Infusi Calumbæ, ℥xi; Aquæ Carui, ℥iv. M. ℥ii bis die.—*In dyspepsia. Child 1 to 2 years old.*
- ℞ Acidi Phosphorici diluti, ℥iss; Tincturæ Aconiti, ℥ss; Tinct. Cinchonæ Co., ℥j; Infusum Aurantii ad ℥vi. M. ℥j ter in die.—*In debility.*
- ℞ Acidi Hydrochlorici diluti, ℥iss; Acidi Hydrocyanici diluti, ℥xx; Infusum Chiratae ad ℥vi. M. ℥j ter in die.—*In gouty dyspepsia.*
- ℞ Quiniæ Sulphatis, Ferri Sulphatis, ā ā gr. xii; Liq. Strychniæ, ℥xxx; Acidi Sulphurici Aromatici, ℥iss; Infusum Quassiæ ad ℥vi. M. ℥j ter in die.—*In neuralgia, &c.*
- ℞ Tinct. Ferri Perchloridi, ℥ss; Potassæ Chloratis. ℥j; Syr., ℥ss; Aquam ad ℥vi. M. ℥j quartâquâque horâ.—*In diphtheria.*

R Quiniæ Sulphatis, gr. xii ; Tincturæ Ferri Perchloridi, ʒii ; Tincturæ Nucis Vomicae, ʒj ; Tincturæ Lupuli, ʒvi ; Magnesiæ Sulphatis, ʒj ; Infusum Lupuli ad ʒvi. M. A sixth part daily, two or three hours after breakfast.—*In constipation with debility.*

R Tincturæ Nucis Vomicae, ʒj ; Acidi Nitrici diluti, ʒj ; Acidi Hydrochlorici diluti, ʒj ; Tincturæ Aurantii, ʒvi ; Aquam ad ʒvi. M. ʒss ter in die.—*In nervous headache, where strychnia does not agree.*

R Potassii Iodidi ; Ammonię Hydrochloratis ā ā ʒiss ; Inf. Humuli ad ʒvi. M. ʒss ter in die.—*In rheumatic headache.*

R Tinct. Ferri Perchloridi, ʒij ; Spiritus Ætheris Nit., ʒiv ; Inf. Quassię ad ʒvi. M. ʒss ter in die.—*In second stage of nephritis.* (Dr Grainger Stewart.)

R Acidi Hydrochlorici, ʒj ; Tinct. Nucis Vomicae, ʒj ; Inf. Gentianę ad ʒvi. M. ʒj ter in die ante cibo.—*Useful in dyspepsia with acid eructations, if given an hour or so before each meal.*

- ℞ Sodæ Bicarb., ℥iss; Spiritūs Ammoniae Aromatici, ℥ii; Tincturæ Gentianæ Compositæ, ℥ss; Syrupi Aurantii, ℥ss; Infusum Gentianæ Compositum ad ℥viii. M. ℥j ter in die.—*In nervous headache with dyspepsia, furred tongue, and acidity.*
- ℞ Tinct. Ferri Perchloridi, ℥ii; Liquoris Strychniæ, ℥j; Glycerine, ℥ss; Infusum Quassiae ad ℥vi. M. ℥ss ter in die.—*In headache of cerebral anæmia, with nervous exhaustion. (Dr Smith.)*
- ℞ Potassæ Bicarbonatis, ℥j; Tincturæ Nucis Vomicae, ℥j; Inf. Gentianæ ad ℥vi. M. ℥j ter in die.—*In dyspepsia with diminished acid secretion, if given before meals.*
- ℞ Beberiae Sulphatis, ℥ss; Syrupi, ℥j; Acidi Sulphurici dil., ℥xxv; Tincturæ Aurantii, ℥j; Aquæ, ℥iv. M. ℥ss t.d.s. —*Intermittent fevers. (Christison.)*

UTERINE THERAPEUTICS.

- ℞ Tinct. Cannabis Indicæ, ℥xx; Spiritus Juniperi, ℥xxx; Spt. Ætheris, ℥xlv; Tincturæ Aconiti, ℥x; Mucilaginis Acaciæ ad ℥iss. M. Fiat Haustus.—*At bedtime, in neuralgic dysmenorrhœa, &c.*
- ℞ Antimonii Tartarati, gr. ii; Tincturæ Opii, ℥xx; Aquæ, ℥vi M. Sumat ℥ss omni hora donec nausea vel vomitus supervenerit.—*In rigidity of the os.*
- ℞ Acidi Benzoici, gr. x; Collodion ℥iv; Acidi Tannici, gr. xv; Acidi Carbolici, gr. xxv. M. Fiat lotio. To be applied to the affected parts.—*In ulceration of cervix uteri.* (Atthill.)
- ℞ Boracis, ℥ss; Morphicæ Sulphatis, gr. vi; Aquæ Rosæ ℥vii. M. Fiat lotio.—*In pruritus of vulvæ.*
- ℞ Liquoris Plumbi Subacetatis, Tincturæ Opii ā ā ℥j. M. A tea-spoonful in half a pint of water, to be injected into the vagina twice daily.—*In leucorrhœa.*

- ℞ Ferri Valerianatis, gr. xviii; Pilulæ Assafœtidæ Compositæ, gr. xxxvi; Olei Sabinæ, ℥xxvi. M. Divide into twelve pills. Take one three times a day.—*In amenorrhœa with hysteria.*
- ℞ Plumbi Iodidi, gr. lxxx; Extracti Belladonnæ, gr. xxiv—xl; Butyri Coeao, ℥j; Olei Olivæ, ℥ij. M. Melt and divide into eight pessaries. One to be placed every night in the vagina.—*In pelvic cellulitis, ovaritis, &c.*
- ℞ Zinci Oxidi, gr. lx; Extracti Belladonnæ, gr. xl; Butyri Coeao, ℥j; Olei Olivæ, ℥ii. M. Divide into eight pessaries.—*In cancer of the uterus and irritable bladder.*
- ℞ Liquoris Potassæ Permanganitis, ℥ii; Aquæ, Oj. M. *An injection in cancer of the uterus. (West.)*
- ℞ Tincturæ Chloroformi Co., ℥ii; Potassii Bromidi, gr. xl; Decoeti Tanaxaci, ℥j; Infusum Auranti ad ℥vii. M. ℥iss ter in die.—*In dysmenorrhœa.*

LIBRARY

INSTITUTE

PUBLICATION
OF
Medical and Scientific Works.



Messrs **E. & S. LIVINGSTONE,**
Medical & Scientific Publishers,

Are prepared to undertake the Publication
of Medical Works on terms advantageous
to Authors.

The rapid sale of Works hitherto placed
in their hands enables them to guarantee
a favourable reception in the Profession,
among Students, and by the Trade.

Applications for ESTIMATES should be
addressed to—

E. & S. LIVINGSTONE,
15 TEVIOT PLACE, EDINBURGH.

PUBLISHED ANNUALLY,

Price 1s; by post, 1s 3d,

Edinburgh Medical School: Guide to Students.

Contains specimens of the Examination Questions, and also Names of the Gentlemen who have passed the First, Second, and Final Professional Examinations.

"It is full of most useful information. It ought to be a most valuable work of reference for the medical student."—*Edinburgh Medical Journal*.

"Contains much valuable information for those intending to enter the medical profession, and especially for those proposing to study in Edinburgh. The Guide contains the regulations of the University and Colleges of Physicians and Surgeons, together with the full syllabuses of all the lecturers and teachers in the School. Much time and money may be saved by a careful perusal of this little book."—*Medical Press and Circular*.

EDINBURGH :

E. & S. LIVINGSTONE, 15 TEVIOT PLACE.

LONDON, Bailliere, Tyndall, and Cox.

„ Simpkin, Marshall, and Co.

DUBLIN, Fannin and Co.; Gill and Son.

GLASGOW, Stenhouse; Hadden; Sime.

ABERDEEN, J. G. Bisset.

MANCHESTER, ... J. E. Cornish.

Just published, price 9s,

The Student's Handbook of Chemistry,

By H. LEICESTER GREVILLE,

F.I.C., F.C.S., ETC.

THIS book contains a large amount of information, condensed into Tables, and also the most important Chemical Calculations.

"The tabulation of facts, containing groups of elements and compounds, is a good feature in this book, and likely to prove very useful to the student. The chapters dealing with organic chemistry are clear and succinct."—*Nature*.

"This little book has many merits. It is clearly and pleasantly written, is well arranged, and contains a great deal of useful and important fact, much of it embodied in tables."—*Lancet*.

"The calculations are very useful."—*Scotsman*.

"A series of tables forms a useful feature of the work."—*Pharmaceutical Journal*.

EDINBURGH :

E. & S. LIVINGSTONE, 15 TEVIOT PLACE.

LONDON, Bailliere, Tyndall, and Cox.

„ Simpkin, Marshall, and Co.

DUBLIN, Fannin and Co.; Gill and Son.

GLASGOW, Stenhouse; Hadden; Sime.

ABERDEEN, J. G. Bisset.

MANCHESTER, J. E. Cornish.

Second edition, price 12s 6d,

Principles and Practice of Midwifery.

By ALEXANDER MILNE, M.D.,

Ex-Vice-President of the Edinburgh Obstetrical
Society.

"The favourable reception accorded to the first edition was well merited, the work being thoroughly practical, and very readable."—*Students' Journal*.

"Dr Milne has contrived to condense a great amount of obstetric doctrine, both physiological and pathological, into his book."—*Lancet*.

"I estimate highly the thought, the labour, and the courage expended in the work."—*Dr J. Matthews Duncan, London*.

"A very excellent and comprehensive manual."—*Professor Graily Hewitt, London*.

"An admirable digest of the subject."—*Professor A. Simpson, Edinburgh*.

EDINBURGH :

E. & S. LIVINGSTONE, 15 TEVIOT PLACE.

LONDON, Bailliere, Tyndall, and Cox.

„ Simpkin, Marshall, and Co.

DUBLIN, Fannin and Co.; Gill and Son.

GLASGOW, Stenhouse ; Hadden ; Sime.

ABERDEEN, J. G. Bisset.

MANCHESTER, ... J. E. Cornish.

Price 10s 6d,

THE STUDENT'S HANDBOOK
OF
Forensic Medicine and Medical Police,

By H. AUBREY HUSBAND, M.B.,

C. M., F. R. C. S. E.,

Lecturer on Medical Jurisprudence and Public
Health in the Extra-Academical School,
Edinburgh, &c.

"The arrangement of the matter is simple, concise, and accurate, and with this book within their reach students should have no excuse for neglecting one of the most important branches of their professional education."—*Medical Press and Circular*.

"The author of this little work has read extensively, both in the domain of medicine and in the region just outside of it, and this extended knowledge, and the happy union of purely medical facts with others of more general interest, give the book an individuality, and certainly have the effect of making it very readable in spite of the great compression of its contents."—*Lancet*.

EDINBURGH :

E. & S. LIVINGSTONE, 15 TEVIOT PLACE.

LONDON, Bailliere, Tyndall, and Cox.

" Simpkin, Marshall, and Co.

DUBLIN, Fannin and Co.; Gill and Son.

GLASGOW, Stenhouse; Hadden; Sime.

ABERDEEN, J. G. Bisset.

MANCHESTER, . . . J. E. Cornish.

Just published, 3rd edition, price 7s 6d,

THE STUDENT'S
Handbook of the Practice of Medicine,

By H. AUBREY HUSBAND, M.B.,
M.C., F.R.C.S.E.,

Lecturer on Medical Jurisprudence and Public
Health in the Extra-Academical School,
Edinburgh, &c.

"The book before us is an excellent example of its class; the whole of medicine is digested and tabulated, and, in fact, served up ready prepared at infinite pains to render it easily assimilated."—*Lancet*.

"The distinguishing features of Dr Husband's Handbook are condensation and clearness of diction—qualities which are certain to ensure for the book a favourable reception by those for whom it is intended. We have pleasure in recommending it to our readers as a most useful work to peruse immediately before presenting one's-self for a medical examination."—*Students' Journal*.

EDINBURGH :

E. & S. LIVINGSTONE, 15 TEVIOT PLACE.

LONDON, Bailliere, Tyndall, and Cox.

„ Simpkin, Marshall, and Co.

DUBLIN, Fannin and Co.; Gill and Son.

GLASGOW, Stenhouse; Hadden; Sime.

ABERDEEN, J. G. Bisset.

MANCHESTER, ... J. E. Cornish.

Price One Shilling.

REFERENCES TO NERVES:

Being Tabular views of the Motor Nerves
to Muscles, etc.

Invaluable to the Anatomical Student.

Cloth, price One Shilling,

POSOLOGICAL TABLES;

Also, Appendix on Poisons:

Being a Tabular arrangement of all the Medicines
contained in the British Pharmacopœia, with Dose,
etc.

TO BOTANICAL STUDENTS.

Price One Shilling,

Thirty Selected Natural Orders of Plants

According to the Programme of the University
of Edinburgh, with Appendices.

Cloth, price 1s. 6d.,

SYLLABUS of VERTEBRATE ZOOLOGY

By J. E. WEST.

This is a most important little book for Students.

Price 4s. 6d.,

INTERSTITIAL NEPHRITES;

Being an Investigation into its Microscopical
Anatomy. With Illustrations.

By BRYAN CHARLES WALLER, M.D.,
F.R.C.S.E.

E. & S. LIVINGSTONE, 15 TEVIOT PLACE.

Price 9s,

Manual of Materia Medica,

By WILLIAM CRAIG, M.D.,

M.C., F.R.S.E.,

Lecturer on Materia Medica, Edinburgh School of
Medicine.

"As a compendium of Materia Medica and Therapeutics, of moderate size but full of information, we know few to surpass, even if to equal, the Manual before us."—*Students' Journal*.

"This book contains sound sense and original information, and will prove a good pocket companion for the student."—*Medical Times and Gazette*.

"Your Manual will be useful to me in various ways, teaching included, and I should think it will be regarded as a boon by students preparing for their examination."—*Professor Harvey, Aberdeen*.

EDINBURGH :

E. & S. LIVINGSTONE, 15 TEVIOT PLACE.

LONDON, Bailliere, Tyndall, and Cox.

„ Simpkin, Marshall, and Co.

DUBLIN, Fannin and Co.; Gill and Son.

GLASGOW, Stenhouse ; Hadden ; Sime.

ABERDEEN, J. G. Bisset.

MANCHESTER, ... J. E. Cornish.

